



Second Semester B.E. Degree Examination, Dec.2013/Jan.2014
Engineering Mathematics – II

Time: 3 hrs.

Max. Marks:100

- Note: 1. Answer any FIVE full questions, choosing at least two from each part.**
2. Answer all objective type questions only on OMR sheet page 5 of the answer booklet.
3. Answer to objective type questions on sheets other than OMR will not be valued.

PART – A

- 1 a.** Choose the correct answers for the following : (04 Marks)
- i) Suppose the equation to be solved is of the form, $y = f(x, \phi)$ then differentiating x we get equation of the form,
 A) $\phi\left(x, p, \frac{dp}{dy}\right) = 0$ B) $\phi\left(y, p, \frac{dp}{dx}\right) = 0$ C) $\phi(x, y, p) = 0$ D) $\phi(x, y, 0) = 0$
- ii) The general solution of the equation, $p^2 - 3p + 2 = 0$ is,
 A) $(y + x - c)(y + 2x - c)$ B) $(y - x - c)(y - 2x - c) = 0$
 C) $(-y - x - c)(y - 2x - c) = 0$ D) $(y - x - c)(y + x - c) = 0$
- iii) Clairaut's equation is of the form,
 A) $x = py + f(p)$ B) $y = p^2 + f(p)$ C) $y = px + f(p)$ D) None of these
- iv) Singular solution of $y = px + 2p^2$ is,
 A) $y^2 + 8y = 0$ B) $x^2 - 8y = 0$ C) $x^2 + 8y - c = 0$ D) $x^2 + 8y = 0$
- b.** Solve $p^2 + 2p \cosh x + 1 = 0$. (04 Marks)
- c.** Find singular solution of $p = \sin(y - xp)$. (06 Marks)
- d.** Solve the equation, $y^2(y - xp) = x^4 p^2$ using substitution $X = \frac{1}{x}$ and $Y = \frac{1}{y}$. (06 Marks)
- 2 a.** Choose the correct answers for the following : (04 Marks)
- i) A second order linear differential equation has,
 A) two arbitrary solution B) One arbitrary solution
 C) no arbitrary solution D) None of these
- ii) If $2, 4i$ and $-4i$ are the roots of A.E of a homogeneous linear differential equation then its solution is,
 A) $e^x + e^x(\cos 4x + \sin 4x)$ B) $C_1 e^{2x} + C_2 \cos 4x + C_3 \sin 4x$
 C) $C_1 e^{2x} + C_2 e^x \cos 4x + C_3 e^x \sin 4x$ D) $C_1 e^{2x} \cos 4x + C_2 e^{2x} \sin 4x$
- iii) P.I. of $(D+1)^2 y = e^{-x+3}$
 A) $\frac{x^2}{2}$ B) $x^3 e^x$ C) $\frac{x^3}{3} e^{-x+3}$ D) $\frac{x^2}{2} e^{-x+3}$
- iv) Particular integral of $f(D)y = e^{ax} V(x)$ is,
 A) $\frac{e^{ax} V(x)}{f(D)}$ B) $e^{ax} \frac{1}{f(D)} [V(x)]$ C) $e^{ax} \frac{1}{f(D+a)} [V(x)]$ D) $\frac{1}{f(D+a)} [e^{ax} V(x)]$
- b.** Solve $\frac{d^3 y}{dx^3} - 3 \frac{d^2 y}{dx^2} + 3 \frac{dy}{dx} - y = 0$. (04 Marks)
- c.** Solve $y'' - 3y' + 2y = 2 \sin x \cos x$ (06 Marks)
- d.** Solve the system of equation, $\frac{dx}{dt} - 2y = \cos 2t$, $\frac{dy}{dt} + 2x = \sin 2t$. (06 Marks)

- 3 a. Choose the correct answers for the following : (04 Marks)
- i) In $x^2y'' + xy' - y = 0$ if $e^t = x$ then we get x^2y'' as,
 A) $(D-1)y$ B) $(D+1)y$ C) $D(D+1)y$ D) None of these
- ii) In second order homogeneous differential equation, $P_0(x)y'' + P_1(x)y' + P_2(x)y = 0$, $x = a$ is a singular point if,
 A) $P_0(a) > 0$ B) $P_0(a) \neq 0$ C) $P_0(a) = 0$ D) $P_0(a) < 0$
- iii) The general solution of $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} - y = 0$ is,
 A) $y = C_1x - C_2 \frac{1}{x}$ B) $C_1x + C_2 \frac{1}{x}$ C) $C_1x + C_2x$ D) $C_1x - C_2x$
- iv) Frobenius series solution of second order linear differential equation is of the form,
 A) $x^m \sum_{r=0}^{\infty} a_r x^r$ B) $\sum_{r=0}^{\infty} a_r x^r$ C) $\sum_{r=0}^{\infty} a_r x^{m+r}$ D) None of these
- b. Solve $y'' + a^2y = \sec ax$ by the method of variation of parameters. (04 Marks)
- c. Solve $x^2 \frac{d^2y}{dx^2} + 4x \frac{dy}{dx} + 2y = e^x$. (06 Marks)
- d. Obtain the series solution of $\frac{dy}{dx} - 2xy = 0$. (06 Marks)
- 4 a. Choose the correct answers for the following : (04 Marks)
- i) PDE of $az + b = a^2x + y$ is,
 A) $\frac{\partial z}{\partial x} \cdot \frac{\partial z}{\partial y} = 1$ B) $\frac{\partial z}{\partial x} \cdot \frac{\partial z}{\partial y} = 0$ C) $\frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} = 0$ D) $\frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} = 1$
- ii) The solution of PDE $Z_{xx} = 2y^2$ is,
 A) $z = x^2 + xf(y) + g(y)$ B) $z = x^2y^2 + xf(y) + g(y)$
 C) $z = x^2y^2 + f(x) + g(x)$ D) $z = y^2 + xf(y) + g(y)$
- iii) The subsidiary equations of $(y^2 + z^2)p + x(yq - z) = 0$ are,
 A) $\frac{dx}{p} = \frac{dy}{q} = \frac{dz}{R}$ B) $\frac{dx}{y^2 + z^2} = \frac{dy}{x} = \frac{dz}{xz}$
 C) $\frac{dx}{y^2 + z^2} = \frac{dy}{xy} = \frac{dz}{xz}$ D) None of these
- iv) In the method of separation of variables to solve $xz_n + z_t = 0$ the assumed solution is of the form,
 A) $X(x)Y(y)$ B) $X(y)Y(y)$ C) $X(t)Y(t)$ D) $X(x)T(t)$
- b. Solve $\frac{\partial^3 z}{\partial x^2 \partial y} = \cos(2x + 3y)$. (04 Marks)
- c. Solve $xp - yq = y^2 - x^2$ (06 Marks)
- d. Solve $3u_x + 2u_y = 0$ by the separation of variable method given that $u = 4e^{-x}$ when $y = 0$. (06 Marks)

PART – B

- 5 a. Choose the correct answers for the following : (04 Marks)
- i) $\int_0^1 \int_0^{x^2} e^{\sqrt{y}} dy dx = \underline{\hspace{2cm}}$
 A) 1 B) - 1/2 C) 1/2 D) None of these
- ii) The integral $\iint_R f(x,y) dx dy$ by changing to polar form becomes,
 A) $\iint_R \phi(r,\theta) dr d\theta$ B) $\iint_R f(r,\theta) dr d\theta$ C) $\iint_R f(r,\theta) r dr d\theta$ D) $\iint_R \phi(r,\theta) r dr d\theta$
- iii) For a real positive number n. the Gamma function $\Gamma(n) = \underline{\hspace{2cm}}$
 A) $\int_0^{\infty} x^{n-1} e^{-x} dx$ B) $\int_0^1 x^{n-1} e^{-x} dx$ C) $\int_0^x x^n e^{-x} dx$ D) $\int_0^1 x^n e^{-x} dx$
- iv) The Beta and Gamma functions relation for $B(m,n) = \underline{\hspace{2cm}}$
 A) $\frac{\Gamma(m)\Gamma(n)}{\Gamma(m-n)}$ B) $\frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}$ C) $\Gamma(m)\Gamma(n)$ D) $\Gamma(mn)$
- b. By changing the order of integration evaluate, $\int_0^a \int_{x/a}^{\sqrt{x/a}} (x^2 + y^2) dy dx$, $a > 0$. (04 Marks)
- c. Evaluate $\int_0^a \int_0^{x+y} \int_0^{x+y+z} e^{x+y+z} dz dy dx$. (06 Marks)
- d. Express the integral $\int_0^1 \frac{dx}{\sqrt{1-x^n}}$ in terms of the Gamma function. Hence evaluate $\int_0^1 \frac{dx}{\sqrt{1-x^{2/3}}}$. (06 Marks)
- 6 a. Choose the correct answers for the following : (04 Marks)
- i) The scalar surface integral of \vec{f} over s, where s is a surface in a three-dimensional region R is given by, $\int \vec{f} \cdot \vec{n} ds = \underline{\hspace{2cm}}$ by Gauss divergence theorem.
 A) $\iiint_V \nabla \cdot \vec{f} dV$ B) $\iint_S \nabla \cdot \vec{t} dx dy$ C) $\iiint_V \nabla \cdot \vec{F} dV$ D) None of these
- ii) If all the surface are closed in a region containing volume V then the following theorem is applicable.
 A) Stoke's theorem B) Green's theorem C) Gauss divergence theorem D) None of these
- iii) The value of $\int_C \{(2xy - x^2) dx + (x^2 + y^2) dy\}$ by using Green's theorem is,
 A) Zero B) One C) Two D) Three
- iv) $\iiint_S f \cdot \vec{n} ds = \underline{\hspace{2cm}}$, where $f = xi+yj+2k$ and S is the surface of the sphere $x^2 + y^2 + z^2 = a^2$
 A) $4\pi a$ B) $4\pi a^2$ C) $4\pi a^3$ D) 4π
- b. Find the work done by a force $f = (2y - x^2)i + 6yzj - 8xz^2k$ from the point (0, 0, 0) to the point (1, 1, 1) along the straight-line joining these points. (04 Marks)
- c. If C is a simple closed curve in the xy-plane, prove by using Green's theorem that the integral $\int_C \frac{1}{2}(x dy - y dx)$ represents the area A enclosed by C. Hence evaluate $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$. (06 Marks)
- d. Verify Stoke's theorem for $\vec{f} = (2x - y)i - yz^2j - y^2zk$ for the upper half of the sphere $x^2 + y^2 + z^2 = 1$. (06 Marks)

7 a. Choose the correct answers for the following : (04 Marks)

i) $L[t^n] = \underline{\hspace{2cm}}$

A) $\frac{n}{s^{n+1}}$ B) $\frac{n}{s^{n-1}}$ C) $\frac{n!}{s^{n-1}}$ D) $\frac{n!}{s^{n+1}}$

ii) $L[e^{-3t}] = \underline{\hspace{2cm}}$

A) $\frac{3}{s-3}$ B) $\frac{3}{s+3}$ C) $\frac{1}{s+3}$ D) $\frac{1}{s-3}$

iii) $L\{f(t-a)H(t-a)\}$ is equal to,

A) $\frac{3!}{(s+2)^4}$ B) $\frac{3!}{(s-2)^4}$ C) $\frac{3}{(s-2)^4}$ D) $\frac{3}{(s-2)}$

iv) $L\{\delta(t-1)\} = \underline{\hspace{2cm}}$

A) e^{-s} B) e^s C) e^{as} D) e^{-as}

b. Evaluate $L\{\sin^3 2t\}$. (06 Marks)

c. Find $L\{f(t)\}$, given that $f(t) = \begin{cases} 2, & 0 < t < 3 \\ t, & t > 3 \end{cases}$. (06 Marks)

d. Express $f(t) = \begin{cases} t^2 & 0 < t < 2 \\ 4t & 2 < t \leq 4 \\ 8 & t > 4 \end{cases}$ in terms of unit step function and hence find their Laplace transform. (04 Marks)

8 a. Choose the correct answers for the following : (04 Marks)

i) $L^{-1}\{\cos at\} = \underline{\hspace{2cm}}$

A) $\frac{s}{s^2 + a^2}$ B) $\frac{s}{s^2 - a^2}$ C) $\frac{1}{s^2 + a^2}$ D) $\frac{1}{s^2 - a^2}$

ii) $L^{-1}\{f(s-a)\} = \underline{\hspace{2cm}}$

A) $e^t f(t)$ B) $e^{at} f(t)$ C) $e^{-at} f(t)$ D) None of these

iii) $L^{-1}\left\{\cot^{-1}\left(\frac{2}{s}\right)\right\} = \underline{\hspace{2cm}}$

A) $\frac{\sin t}{t}$ B) $\frac{\sinh at}{t}$ C) $\frac{\sin at}{t}$ D) $\frac{\sinh t}{t}$

iv) For the function $f(t) = 1$, convolution theorem condition,

A) Not satisfied B) Satisfied with some condition
C) Satisfied D) None of these

b. Find the inverse Laplace transform of $\frac{2s^2 - 6s + 5}{(s-1)(s-2)(s-3)}$. (04 Marks)

c. Find $L^{-1}\left\{\frac{s}{(s-1)(s^2+4)}\right\}$ using convolution theorem. (06 Marks)

d. Solve differential equation $y''(t) + y = F(t)$, where $F(t) = \begin{cases} 0 & 0 < t < 1 \\ 2 & t > 1 \end{cases}$. Given that $y(0) = 0 = y'(0)$. (06 Marks)

* * * * *