

USN

|  |  |  |  |  |  |  |  |  |  |
|--|--|--|--|--|--|--|--|--|--|
|  |  |  |  |  |  |  |  |  |  |
|--|--|--|--|--|--|--|--|--|--|

10MAT21

**Second Semester B.E. Degree Examination, June/July 2018**  
**Engineering Mathematics – II**

Time: 3 hrs.

Max. Marks:100

**Note: Answer any FIVE full questions, choosing at least two from each part.**

**PART – A**

- 1 a. Choose the correct answers for the following : (04 Marks)

i) If the differential equation is solving for x then it is of the form,

A)  $x = f\left(\frac{P}{y}\right)$       B)  $y = f(x, P)$       C)  $x = f\left(\frac{y}{P}\right)$       D)  $x = f(y, p)$

ii) The general solution of  $P^2 - 7P + 12 = 0$ ,

A)  $(y + 3x - c)(y + 4x - c) = 0$       B)  $(y - 3x - c)(y - 4x - c) = 0$   
 C)  $(y - 4x)(y + 3x) = 0$       D) None of these

iii)  $y = cx + f(c)$  is the general solution of the equation,

A)  $x = py + f(p)$       B)  $y = 3x + \log p$       C)  $y = px + f(p)$       D) None of these

iv) The general solution of the equation,  $(y - px)^2 = 4p^2 + 9$  is,

A)  $y = Cx + \sqrt{4C^2 + 9}$       B)  $y = C + \sqrt{4C^2 + 9}$   
 C)  $y = Cx + \sqrt{4C^2 - 9}$       D)  $y - Cx = 4C^2 + 9$

b. Solve  $P^2 + 2Py \cot x = y^2$ .

c. Solve  $(y - px)(p - 1) = p$ .

d. Solve the equation  $(px - y)(py + x) = 2p$  by reducing into Clairaut's form, taking the substitutions  $X = x^2$ ,  $Y = y^2$ . (06 Marks)

- 2 a. Choose the correct answers for the following : (04 Marks)

i)  $\frac{1}{f(D)} [e^{2x} \cdot x^2] = \underline{\hspace{2cm}}$ ,

A)  $e^{2x} \frac{1}{f(D+2)} x^2$       B)  $e^{2x} \frac{1}{f(D-2)} x^2$       C)  $x^2 \frac{1}{f(D-2)} e^{2x}$       D)  $x^2 \frac{1}{f(D+2)} e^{2x}$

ii) If 1, 1, -2 are the roots of auxiliary equation of the differential equation then its solution is,

A)  $e^x + e^x + e^{-2x}$       B)  $(C_1 + C_2 x)e^x + C_3 e^{-2x}$   
 C)  $C_1 e^x + C_2 e^x + C_3 e^{-2x}$       D) None of these

iii) Particular integral of  $(D+1)^2 y = e^{-x+3}$  is,

A)  $\frac{x^2}{2}$       B)  $x^3 e^x$       C)  $\frac{x^3}{3} e^{-x+3}$       D)  $-\frac{x^2}{2} e^{-x+3}$

iv) General solution of non homogeneous linear differential equation is,

- A) Sum of complementary function and particular integral  
 B) Difference of complementary function and particular integral  
 C) Product of complementary function and particular integral  
 D) None of these

b. Solve:  $\frac{d^3 y}{dx^3} + 3 \frac{d^2 y}{dx^2} + 3 \frac{dy}{dx} - y = 0$ . (05 Marks)

c. Solve:  $\frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 2y = 2 \sin x \cos x$ . (05 Marks)

d. Solve the system of equations:  $\frac{dx}{dt} - 2y = \cos 2t$ ,  $\frac{dy}{dt} + 2x = \sin 2t$ . (06 Marks)

3 a. Choose the correct answers for the following :

i) The complementary function of  $x^2 \frac{d^2y}{dx^2} + 4x \frac{dy}{dx} + 2y = e^x$

- A)  $C_1 e^{-x} + C_2 e^{-3x}$     B)  $C_1(-x) + C_2(-2x)$     C)  $C_1 e^{-2x} + C_2 e^{2x}$     D)  $\frac{C_1}{x} + \frac{C_2}{x^2}$

ii) In the method of variation of parameters the value of W is called \_\_\_\_\_,

- A) Wronskian function    B) Exponential function  
C) DeMorgans function    D) Euler's function

iii) The roots of the auxillary equation of the transformed equation of,

$$(2x+1)^2 \frac{d^2y}{dx^2} - 2(2x+1) \frac{dy}{dx} - 12y = 6x + 5 \text{ are,}$$

- A) 3, -1    B) -3, 1    C) 12, -4    D) None of these

iv) The solution of  $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} = 0$  is,

- A)  $y = C_1 \cos x + C_2 \sin x$     B)  $y = C_1 e^x + C_2 e^{-x}$     C)  $y = C_1 \log x + C_2$     D)  $y = C_1 + 6x$

b. Solve  $y'' - 6y' + 9y = \frac{e^{3x}}{x^2}$  by the method of variation of parameters. (05 Marks)

c. Solve  $x^2 \frac{d^2y}{dx^2} - 3x \frac{dy}{dx} + 4y = \sin(\log x)$ . (05 Marks)

d. Solve  $(1+x^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx} - y = 0$  in series solution. (06 Marks)

4 a. Choose the correct answers for the following :

i) The partial differential equation obtained by eliminating arbitrary constants from the relation  $z = (x-a)^2 + (y-b)^2$  is,

- A)  $p^2 + q^2 = 4z$     B)  $p^2 - q^2 = 4z$     C)  $p + q = z$     D)  $p - q = 2z$

ii) The auxillary equation of  $Pp + Qq = R$  are,

- A)  $\frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R}$     B)  $\frac{dx}{p} = \frac{dy}{q} = \frac{dz}{R}$     C)  $\frac{dx}{x} = \frac{dy}{y} = \frac{dz}{z}$     D)  $\frac{dx}{x} + \frac{dy}{y} + \frac{dz}{z} = 0$

iii) A solution of  $(y-z)p + (z-x)q = x-y$  is . . .

- A)  $x^2 - y^2 - z^2 = f(x-y+z)$     B)  $x^2 + y^2 + z^2 = f(x+y+z)$   
C)  $x^2 - y^2 - z^2 = f(x-y-z)$     D) None of these

iv) By the separation of variables, solution is in the form of,

- A)  $Z = X.Y$     B)  $Z = X+Y$     C)  $Z = X^2.Y^2$     D)  $Z = \frac{X}{Y}$

b. Form a PDE from the relation  $Z = y^2 + 2f\left[\frac{1}{x} + \log y\right]$ . (05 Marks)

c. Solve the equation  $(x^2 - y^2 - z^2)p + 2xyq = 2xz$ . (05 Marks)

d. Use the method of separation of variables to solve  $\frac{\partial u}{\partial x} = 2 \frac{\partial u}{\partial t} + u$ , given that  $u(x,0) = 6e^{-x}$ . (06 Marks)

### PART - B

5 a. Choose the correct answers for the following :

(04 Marks)

i) The value of  $\int_1^2 \int_1^3 xy^2 dx dy$  is \_\_\_\_\_,

- A) 0    B) 1    C) 13/2    D) 13

ii) The integral  $\int_0^1 \int_{\sqrt{y}}^1 dx dy$  after changing the order of integration,

A)  $\int_0^2 \int_0^y dx dy$

B)  $\int_0^1 \int_0^{y^2} dx dy$

C)  $\int_0^1 \int_0^1 dx dy$

D)  $\int_0^1 \int_0^{x^2} dy dx$

iii)  $\beta\left(\frac{1}{2}, \frac{1}{2}\right) = \underline{\hspace{2cm}}$

A)  $\sqrt{\pi}$

B)  $\frac{2}{\sqrt{\pi}}$

C)  $\pi\sqrt{2}$

D) 3.1416

iv) The value of  $\Gamma(1/4)\Gamma(3/4) = \underline{\hspace{2cm}}$

A)  $2\sqrt{\pi}$

B)  $\pi\sqrt{2}$

C)  $\sqrt{2}$

D)  $\frac{\sqrt{\pi}}{2}$

b. Change the order of integration in  $\int_0^1 \int_0^{\sqrt{1-x^2}} y^2 dx dy$  and hence evaluate the same. (05 Marks)

c. Show that  $\beta(m, n) = \int_0^{\infty} \frac{x^{m-1}}{(1+x)^{m+n}} dx$ . (05 Marks)

d. Show that  $\int_0^{\pi/2} \frac{d\theta}{\sqrt{\sin \theta}} \times \int_0^{\pi/2} \sqrt{\sin \theta} d\theta = \pi$ . (06 Marks)

6 a. Choose the correct answers for the following : (04 Marks)

i) If  $\int_C \vec{F} \cdot d\vec{r} = 0$  then  $\vec{F}$  is called,

A) Rotational

B) Solenoidal

C) Irrotational

D) Dependent

ii) If  $f$  is a vector field over a region of volume  $V$  in three dimensional space then,  $\int_V f \cdot dv$  is called,

A) Scalar volume integral

B) Vector volume integral

C) Scalar surface integral

D) Vector surface integral

iii) In Green's theorem in the plane  $\iint_A \left[ \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right] dx dy$  is \_\_\_\_\_,

A)  $\int_C [Mdx - Ndy]$

B)  $\int_C [Mdx + Ndy]$

C)  $\int_C [Mdx \times Ndy]$

D)  $\int_C [Ndx - Mdy]$

iv) If  $V$  is the volume bounded by a surface  $S$  and  $\vec{F}$  is a continuously differentiable vector function then,  $\iiint_V \text{div } \vec{F} dv = \underline{\hspace{2cm}}$

A) 0

B)  $\iint_S \vec{F} \times \hat{n} ds$

C)  $\iint_S \vec{F} \cdot \hat{n} ds$

D) None of these

b. Evaluate  $\iint_S f \cdot \hat{n} ds$  where  $f = yz\hat{i} + zx\hat{j} + xy\hat{k}$  and  $S$  is the surface of the sphere  $x^2 + y^2 + z^2 = a^2$  in the 1<sup>st</sup> Octant. (05 Marks)

c. Verify Green's theorem for  $\int_C (xy + y^2) dx + x^2 dy$  where  $C$  is the closed curve made up of the line  $y = x$  and the parabola  $y = x^2$ . (05 Marks)

d. Verify stoke's theorem for,  $f = (2xy - y)\hat{i} - yz^2\hat{j} - y^2z\hat{k}$  for the upper half of the sphere  $x^2 + y^2 + z^2 = 1$  (06 Marks)

7 a. Choose the correct answers for the following :

i) If  $L[f(t)] = \tilde{f}(s)$  then  $L[e^{at}f(t)]$  is \_\_\_\_\_,  
 A)  $\tilde{f}(s-a)$       B)  $\tilde{f}(s+a)$       C)  $\tilde{f}\left(\frac{s}{a}\right)$       D) None of these

ii)  $L\left[\frac{\sin at}{t}\right] =$  \_\_\_\_\_  
 A)  $\cot^{-1}\left(\frac{s}{a}\right)$       B)  $\tan^{-1}\left(\frac{s}{a}\right)$       C)  $\frac{\pi}{2} + \tan^{-1}\left(\frac{s}{a}\right)$       D) None of these

iii)  $L[u(t+2)] =$  \_\_\_\_\_  
 A)  $\frac{e^{-2s}}{s^2}$       B)  $e^{2s}$       C)  $\frac{e^{2s}}{s}$       D)  $\frac{e^{-2s}}{s}$

iv)  $L[s(t+2)] =$  \_\_\_\_\_  
 A)  $e^{-as}$       B)  $e^{2s}$       C)  $e^{-2s}$       D)  $e^{as}$

b. Find the value of  $\int_0^{\infty} t^3 e^{-t} \sin t dt$  using Laplace Transforms. (05 Marks)

c. If  $f(t) = \begin{cases} t, & 0 \leq t \leq a \\ 2a-t, & a \leq t \leq 2a \end{cases}$  where  $f(t+2a) = f(t)$  show that  $L[f(t)] = \frac{1}{s^2} \tanh\left(\frac{as}{2}\right)$ . (05 Marks)

d. Express  $f(t) = \begin{cases} 1, & 0 < t \leq 1 \\ t, & 1 < t \leq 2 \\ t^2, & t > 2 \end{cases}$  in terms of unit step function and hence find its Laplace transform. (06 Marks)

8 a. Choose the correct answers for the following :

(04 Marks)

i)  $L^{-1}[\cos at] =$  \_\_\_\_\_  
 A)  $\frac{s}{s^2 + a^2}$       B)  $\frac{s}{s^2 - a^2}$       C)  $\frac{1}{s^2 - a^2}$       D)  $\frac{1}{s^2 + a^2}$

ii)  $L^{-1}[\tilde{f}(s-a)] =$  \_\_\_\_\_  
 A)  $e^{at}f(t)$       B)  $e^f(t)$       C)  $e^{-at}f(t)$       D) None of these

iii)  $L^{-1}[\cot^{-1}(s)] =$  \_\_\_\_\_  
 A)  $\frac{\sinh t}{t}$       B)  $\frac{\sin t}{t}$       C)  $-\frac{\sinh t}{t}$       D)  $-\frac{\sin t}{t}$

iv)  $L\left[\int_0^t f(u)g(t-u)du\right] =$  \_\_\_\_\_  
 A)  $f(s)g(s)$       B)  $\frac{f(s)}{g(s)}$       C)  $f(t) \cdot g(t)$       D)  $f(s) + g(s)$

b. Find  $L^{-1}\left[\log \frac{s+1}{s-1}\right]$  (05 Marks)

c. Find  $L^{-1}\left[\frac{s}{(s-1)(s^2+4)}\right]$  using convolution theorem. (05 Marks)

d. Solve the initial value problem  $y'' - 3y' + 2y = 1 - e^{2t}$ ,  $y(0) = 1$ ,  $y'(0) = 1$  using Laplace transform. (06 Marks)

\*\*\*\*\*