

| | | | | | | | | | |
|--|--|--|--|--|--|--|--|--|--|
| | | | | | | | | | |
|--|--|--|--|--|--|--|--|--|--|

First Semester B.E. Degree Examination, December 2011
Engineering Mathematics - I

Time: 3 hrs.

Max. Marks:100

- Note:** 1. Answer any FIVE full questions, choosing at least two from each part.
 2. Answer all objective type questions only on OMR sheet page 5 of the answer booklet.
 3. Answer to objective type questions on sheets other than OMR will not be valued.

PART - A

- 1 a. Choose your answers for the following : (04 Marks)

i) If $y = \frac{x}{x-1}$, then y_n is

A) $\frac{(-1)^{n-1}n!}{(x-1)^{n+1}}$ B) $\frac{(-1)^n n!}{(x-1)^{n+1}}$ C) $\frac{(-1)^n (n+1)!}{(x-1)^{n+1}}$ D) $\frac{(-1)^n n!}{(x-1)^n}$

ii) If $y = \log(ax+b)$, then y_n is

A) $\frac{(-1)^n n! a^n}{(ax+b)^n}$ B) $\frac{(-1)^{n-1} n! a^n}{(ax+b)^{n+1}}$ C) $\frac{(-1)^{n-1} (n-1)! a^n}{(ax+b)^n}$ D) $\frac{(-1)^n (n-1)! a^n}{(ax+b)^{n+1}}$

iii) If $f(x) = \sin x$, $x \in (0, \pi)$, then by Rolle's theorem the value of 'x', where the Tangent is parallel to x - axis.

A) 0 B) $\frac{\pi}{2}$ C) $\frac{\pi}{3}$ D) $\frac{\pi}{4}$

iv) Expansion of $\log(1+x)$ in powers of x is

A) $x + \frac{x^2}{2} + \frac{x^3}{3} + \dots$ B) $x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$
 C) $1 - \frac{x}{1!} + \frac{x^2}{2!} - \frac{x^3}{3!} + \dots$ D) $\frac{x}{1!} - \frac{x^2}{2!} + \frac{x^3}{3!} - \frac{x^4}{4!} + \dots$

- a. If $x = \tan(\log y)$, show that $(1+x^2)y_{n+1} + (2nx-1)y_n + n(n-1)y_{n-1} = 0$. (04 Marks)
 b. State and prove Cauchy's mean value theorem. (06 Marks)
 c. Expand $f(x) = \sin(e^x - 1)$ in power's of 'x' upto the terms containing x^4 . (06 Marks)

- 2 a. Choose your answers for the following : (04 Marks)

i) The indeterminate form of $\lim_{x \rightarrow 1} \left(\frac{x}{x-1} - \frac{(x-1)}{\log x} \right)$ is

A) $\infty - \infty$ B) $\frac{0}{0}$ C) $\frac{\infty}{\infty}$ D) None of these

ii) The angle between the radius vector and the tangent to the curve $r = k e^{\theta \cot \alpha}$, where K and α are constants, is :

A) K B) θ C) α D) 0

iii) The Pedal equation of the curve $r = a\theta$ is.

A) $p^2 = ar$ B) $\frac{1}{p^2} = \frac{a}{r^2}$ C) $\frac{1}{p^2} = \frac{1}{r^2} + a^2$ D) $\frac{1}{p^2} = \frac{1}{r^2} + \frac{a^2}{r^4}$

iv) The radius of curvature at any point 't' on the curve defined by $x = f(t)$, $y = \phi(t)$ is given by

A) $\frac{[(x')^2 + (y')^2]^{3/2}}{x'y'' - y'x''}$ B) $\frac{x'y'' - y'x''}{[(x')^2 + (y')^2]^{3/2}}$ C) $\frac{(x')^2 + (y')^2}{(x'y'' - y'x'')^{3/2}}$ D) $\frac{(x'y'' - y'x'')^{3/2}}{(x')^2 + (y')^2}$

- b. Find the angle of intersection between the curves $r^n \cos(n\theta) = a^n$ and $r^n \sin(n\theta) = b^n$. (04 Marks)
- c. Show that the radius of curvature at any point 'θ' to the curve $x = a(\theta + \sin \theta)$, $y = a(1 - \cos \theta)$, is $4a \cos(\frac{\theta}{2})$. (06 Marks)
- d. Evaluate $\lim_{x \rightarrow 0} \left(\frac{a^x + b^x + c^x}{3} \right)^{\frac{1}{x}}$. (06 Marks)

3 a. Choose your answers for the following : (04 Marks)

- i) If $u = x^{y-1}$, then $\frac{\partial u}{\partial y}$ is
 A) $x^{y-1} \log x$ B) $(y-1)x^{y-2}$ C) $x^{y-1} \log y$ D) $x^y \log x$
- ii) If $Z = f(u, v)$, where $u = x + ct$ and $v = x - ct$, then $\frac{\partial Z}{\partial t}$ is given by
 A) $\frac{\partial z}{\partial u} - \frac{\partial z}{\partial v}$ B) $\frac{\partial z}{\partial u} + \frac{\partial z}{\partial v}$ C) $c \left(\frac{\partial z}{\partial u} - \frac{\partial z}{\partial v} \right)$ D) $c \left(\frac{\partial z}{\partial v} - \frac{\partial z}{\partial u} \right)$
- iii) If $x = u(1-v)$, $y = uv$, then $J \left(\frac{x, y}{u, v} \right)$ is equal to
 A) u B) $\frac{1}{u}$ C) uv D) $\frac{u}{v}$
- iv) The necessary condition for the function $f(x, y)$ to possess extreme values is
 A) $f_x = f_y = 0$ B) $f_{xx} - f_{yy} = 0$ C) $(f_{xx})(f_{yy}) - f_{xy}^2 = 0$ D) $f_x > 0, f_y > 0$

b. If $u = f \left(\frac{y-x}{xy}, \frac{z-x}{xz} \right)$, find $x^2 \frac{\partial u}{\partial x}$. (04 Marks)

c. If $x + y + z = u$, $y + z = v$ and $z = uvw$, show that $J \left(\frac{x, y, z}{u, v, w} \right) = uv$. (06 Marks)

d. The Horse power required to propel a steamer is proportional to the square of the distance and cube of the velocity. If the distance is increased by 4% and velocity increased by 3%, find the percentage of increase in the Horse power. (06 Marks)

4 a. Choose your answers for the following : (04 Marks)

- i) If $\vec{R} = xi + yj + zk$, $|\vec{R}| = r$, then ∇r^2 is equal to
 A) $\frac{\vec{R}}{r^2}$ B) $\frac{-\vec{R}}{2}$ C) $\frac{\vec{R}}{r}$ D) $2\vec{R}$
- ii) If $\vec{F} = 3x^2i - xyj + (a-3)xzk$ is solenoidal, then 'a' is equal to
 A) 0 B) -2 C) 2 D) 3
- iii) If $\vec{A} = x^2i + y^2j + z^2k$, then $\text{curl } \vec{A}$ is given by
 A) $2xi + 2yj + 2zk$ B) 0 C) $\frac{xi + yj + zk}{2}$ D) $2x + 2y + 2z$
- iv) The scale factors for cylindrical coordinate system (ρ, ϕ, z) are given by
 A) $(\rho, 1, 1)$ B) $(1, \rho, 1)$ C) $(1, 1, \rho)$ D) None of these

b. Prove that $\nabla \cdot \phi \vec{F} = \nabla \phi \cdot \vec{F} + \phi (\nabla \cdot \vec{F})$. (04 Marks)

c. If $\vec{F} = 2xy^3z^4i + 3x^2y^2z^4j + 4x^2y^3z^3k$, find i) $(\nabla \cdot \vec{F})$ ii) $\nabla \times \vec{F}$. (06 Marks)

d. Obtain the expression for $\nabla \cdot \vec{F}$ in orthogonal curvilinear coordinate system (u_1, u_2, u_3) . (06 Marks)

PART – B

- 5 a. Choose your answers for the following : (04 Marks)
- i) Given $\int_0^1 x^n dx = \frac{1}{n+1}$, then $\frac{d^2}{dx^2} \int_0^1 x^n dx$ gives
- A) $\int_0^1 (\log x)^2 x^n dx = \frac{2}{(1+n)^2}$ B) $\int_0^1 (\log x)^2 x^n dx = \frac{2}{(1+n)^3}$
- C) $\int_0^1 (\log x)^n x^n dx = \frac{2}{(1+n)^2}$ D) $\int_0^1 (\log x)^2 x^n dx = \frac{-2}{(1+n)^3}$
- ii) The value of the integral $\int_0^{\pi} \sin^6 x \cos^5 x dx$ is
- A) 0 B) $\frac{8}{693}$ C) $\frac{8\pi}{693}$ D) None of these
- iii) The volume of the solid generated by revolving the curve $r = a(1 + \cos\theta)$ about the line $\theta = 0$ is given by
- A) $\frac{2\pi}{3} a^3 \int_0^{\pi} (1 + \cos\theta)^3 \sin\theta d\theta$ B) $\frac{2\pi}{3} a^3 \int_0^{\pi} (1 + \cos\theta)^3 \cos\theta d\theta$
- C) $\frac{2\pi}{3} a^3 \int_0^{2\pi} (1 + \cos\theta)^3 \sin\theta d\theta$ D) $\frac{4\pi a^3}{3}$
- iv) The entire length of the asteroid $x^{2/3} + y^{2/3} = a^{2/3}$ is
- A) 4a B) 8a C) 6a D) 3a
- b. Obtain the reduction formula of the integral $\int \cos^n x dx$. (04 Marks)
- c. Using Leibnitz rule under differentiation under integral sign, evaluate $\int_0^{\pi} \frac{\log(1+2\cos x)}{\cos x} dx$. (06 Marks)
- d. Find the surface generated by revolving the cycloid $x = a(\theta - \sin\theta)$, $y = a(1 - \cos\theta)$ about its base, (consider one arc in the 1st quadrant). (06 Marks)
- 6 a. Choose your answers for the following : (04 Marks)
- i) The general solution of the differential equation $\frac{dy}{dx} = \sec\left(\frac{y}{x}\right) + \frac{y}{x}$ is
- A) $\tan y/x - \log x = c$ B) $\sin(y/x) - \log x = c$
- C) $\operatorname{Cosec}(y/x) - \log x = c$ D) $\cos(y/x) - \log x = c$
- ii) Integrating factor for the differential equation $\frac{dx}{dy} + \frac{2x}{y} = y^2$ is
- A) y^2 B) e^{x^2} C) e^{2y} D) e^{y^2}
- iii) The general solution of the differential equation $(x - y) dx + (y - x) dy = 0$ is
- A) $\frac{x^2}{2} - y - \frac{y^2}{2} = c$ B) $\frac{x^2}{2} - y + \frac{y^2}{2} = c$ C) $\frac{x^2}{2} - yx + \frac{y^2}{2} = c$ D) None of these
- iv) Given the differential equation of $f(r, \theta, c) = 0$, we get differential equation of orthogonal trajectories by changing $r \frac{d\theta}{dr}$ by
- A) $\frac{1}{r} \frac{dr}{d\theta}$ B) $-r^2 \frac{dr}{d\theta}$ C) $\frac{-1}{r} \frac{dr}{d\theta}$ D) $r \frac{dr}{d\theta}$
- b. Solve $(x^2 - 4xy - 2y^2) dx + (y^2 - 4xy - 2x^2) dy = 0$. (04 Marks)
- c. Solve $(x + 2y^3) \frac{dy}{dx} = y$. (06 Marks)
- d. Find the orthogonal trajectories of the family of curves $\frac{x^2}{a^2} + \frac{y^2}{b^2 + \lambda} = 1$ (' λ ' being the parameter). (06 Marks)

7 a. Choose your answers for the following : (04 Marks)

i) The rank of the matrix $\begin{pmatrix} 6 & 1 & 3 & 8 \\ 4 & 2 & 6 & -1 \\ 10 & 3 & 9 & 7 \\ 16 & 4 & 12 & 15 \end{pmatrix}$ is equal to

A) 2 B) 3 C) 4 D) 1

ii) The exact solution of the system of equations $10x + y + z = 12$, $x + 10y + z = 12$, $x + y + 10z = 12$ by inspection is equal to

A) $[0 \ 0 \ 0]^T$ B) $[1 \ 1 \ 1]^T$ C) $[1 \ 1 \ -1]^T$ D) $[-1 \ -1 \ -1]^T$

iii) If the given system of linear equations in 'n' variables is consistent then the number of linearly independent solution is given by

A) n B) n - 1 C) r - n D) n - r

(Where 'r' stands for rank of co-efficient, matrix).

iv) The trivial solution for the given system of equations

$$qx - y + 4z = 0, \quad 4x - 2y + 3z = 0, \quad 5x + y - 6z = 0 \text{ is}$$

A) (1, 2, 0) B) (0 4 1) C) (0 0 0) D) (1 -5 0)

b. Using elementary row transformations find the rank of the matrix $\begin{pmatrix} 0 & 1 & -3 & -1 \\ 1 & 0 & 1 & 1 \\ 3 & 1 & 0 & 2 \\ 1 & 1 & -2 & 0 \end{pmatrix}$. (04 Marks)

c. Test for consistency and solve the system of equations $x + 4 + 3z = 0$, $x - y + z = 0$, $2x - y + 3z = 0$. (06 Marks)

d. Applying Gauss Jordan method solve $2x + 3y - z = 5$, $4x + 4y - 3z = 3$, $2x - 3y + 2z = 2$. (06 Marks)

8 a. Choose your answers for the following : (04 Marks)

i) The linear transformation $y = Ax$ is regular if

A) $|A| = 0$ B) $|A| = 1$ C) $|A| = -1$ D) $|A| \neq 0$

ii) The transformation $\xi = x \cos \alpha - y \sin \alpha$, $\eta = x \sin \alpha + y \cos \alpha$ is orthogonal then the inverse of the transformation matrix is given by

A) $\begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}$ B) $\begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$ C) $\begin{pmatrix} \sin \alpha & \cos \alpha \\ \cos \alpha & -\sin \alpha \end{pmatrix}$ D) $\begin{pmatrix} -\sin \alpha & \cos \alpha \\ \cos \alpha & \sin \alpha \end{pmatrix}$

iii) The eigen vector 'x' of the matrix 'A' corresponding to eigen value ' λ ' satisfy the equation

A) $AX = \lambda X$ B) $\lambda(A - X) = 0$ C) $XA - \lambda A = 0$ D) $|A - \lambda I|X = 0$

iv) Two square matrices A and B are similar if

A) $A = B$ B) $B = P^{-1}AP$ C) $A^1 = B^1$ D) $A^{-1} = B^{-1}$

b. Show that the transformation given below $y_1 = 2x_1 + x_2 + x_3$, $y_2 = x_1 + x_2 + 2x_3$, $y_3 = x_1 - 2x_3$ is regular and find the inverse transformation. (04 Marks)

c. Find the matrix P which diagonalizes the matrix $A = \begin{bmatrix} -1 & 1 & 2 \\ 0 & -2 & -1 \\ 0 & 0 & -3 \end{bmatrix}$. (06 Marks)

d. Reduce the quadratic form $x_1^2 + 3x_2^2 + 3x_3^2 - 2x_2x_3$ in to canonical form by an appropriate orthogonal transformation which transforms $x_1 \ x_2 \ x_3$ in terms of new variables $y_1 \ y_2 \ y_3$. (06 Marks)
