

## Third Semester B.E. Degree Examination, June/July 2015

## **Engineering Mathematics - III**

Time: 3 hrs. Max. Marks: 100

Note: Answer FIVE full questions, selecting at least TWO questions from each part.

## PART - A

1 a. Expand  $f(x) = x \sin x$  as a Fourier series in the interval  $(-\pi, \pi)$ , Hence deduce the following:

i) 
$$\frac{\pi}{2} = 1 + \frac{2}{1.3} - \frac{2}{3.5} + \frac{2}{5.7}$$
  
ii)  $\frac{\pi - 2}{4} = \frac{1}{1.3} - \frac{1}{3.5} + \frac{1}{5.7} - + \dots$  (07 Marks)

b. Find the half-range Fourier cosine series for the function

$$f(x) = \begin{cases} kx, & 0 \le x \le \frac{\ell}{2} \\ k(\ell - x), & \ell \le x \le \ell \end{cases}$$

Where k is a non-integer positive constant.

(06 Marks)

c. Find the constant term and the first two harmonics in the Fourier series for f(x) given by the following table.

x:	0	π/3	$2\pi/3$	π	$4\pi/3$	$5\pi/3$	2π
F(x):	1.0	1.4	1.9	1.7	1.5	1.2	1.0

(07 Marks)

2 a. Find the Fourier transform of the function  $f(x) = xe^{-a|x|}$  (07 Marks)

b. Find the Fourier sine transforms of the

Functions 
$$f(x) = \begin{cases} \sin x, & 0 < x < a \\ 0, & x \ge a \end{cases}$$
 (06 Marks)

c. Find the inverse Fourier sine Transform of

$$F_x(\alpha) = \frac{1}{\alpha} e^{-a\alpha} \quad a > 0.$$
 (07 Marks)

- 3 a. Find various possible solution of one dimensional wave equation  $\frac{\partial^2 u}{\partial t^2} = C^2 \frac{\partial^2 u}{\partial x^2}$  by separable variable method.
  - b. Obtain solution of heat equation  $\frac{\partial u}{\partial t} = C^2 \frac{\partial^2 u}{\partial t^2}$  subject to condition u(0,t) = 0, u(t,t) = 0, u(x,0) = f(x).
  - c. Solve Laplace equation  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$  subject to condition  $u(0, y) = u(\ell, y) = 0$ , u(x, 0) = 0,

$$u(x, a) = \sin\left(\frac{\pi x}{\ell}\right). \tag{07 Marks}$$

a. The pressure P and volume V of a gas are related by the equation  $PV^r = K$ , where r and K are constants. Fit this equation to the following set of observations (in appropriate units)

<b>P</b> :	0.5	1.0	1.5	2.0	2.5	3.0
V:	1.62	1.00	0.75	0.62	0.52	0.46

(07 Marks)

b. Solve the following LPP by using the Graphical method:

Maximize:  $Z = 3x_1 + 4x_2$ 

Under the constraints  $4x_1 + 2x_2 \le 80$ 

$$2x_1 + 5x_2 \le 180$$

$$x_1, x_2 \ge 0.$$
 (06 Marks)

c. Solve the following using simplex method

Maximize: Z = 2x + 4y, subject to the

Constraint:  $3x + y \le 2z$ ,  $2x + 3y \le 24$ ,  $x \ge 0$ ,  $y \ge 0$ .

(07 Marks)

## PART - B

- a. Using the Regular Falsi method, find a real root (correct to three decimal places) of the equation  $\cos x = 3x - 1$  that lies between 0.5 and 1 (Here, x is in radians). (07 Marks)
  - b. By relaxation method

Solve: 
$$-x + 6y + 27z = 85$$
,  $54x + y + z = 110$ ,  $2x + 15y + 6z = 72$ .

c. Using the power method, find the largest eigen value and corresponding eigen vectors of the

matrix 
$$A = \begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix}$$

taking  $[1, 1, 1]^T$  as the initial eigen vectors. Perform 5 iterations.

(07 Marks)

(06 Marks)

a. From the data given in the following Table; find the number of students who obtained (i) Less than 45 marks ii) between 40 and 45 marks

(1) Bess than 45 marks in between 40 and 45 marks.							
Marks	30 - 40	40 – 50	50 – 60	60 - 70	70 - 80		
No. of Students	31	42	51	35	31		

(07 Marks)

b. Using the Lagrange's formula, find the interpolating polynomial that approximates to the function described by the following table:

(06 Marks)

c. Evaluate  $\int_{1+x^2}^{1} dx$  by using Simpson's  $\left(\frac{3}{8}\right)^{th}$  Rule, dividing the interval into 3 equal parts.

Hence find an approximate value of  $\log \sqrt{2}$ .

(07 Marks)

a. Solve the one – dimensional wave equation  $\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2}$ 

Subject to the boundary conditions u(0, t) = 0, u(1, t) = 0,  $t \ge 0$  and the initial conditions

$$u(x, 0) = \sin \pi x, \quad \frac{\partial u}{\partial t}(x, 0) = 0, \quad 0 < x < 1.$$
 (07 Marks)

- b. Consider the heat equation  $2\frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$  under the following conditions:
  - i)  $u(0, t) = u(4, t) = 0, t \ge 0$
  - ii) u(x, 0) = x(4 x), 0 < x < 4.

Employ the Bendre – Schmidt method with h = 1 to find the solution of the equation for  $0 \le t \le 1$ .

c. Solve the two – dimensional Laplace equation  $\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial y^2} = 0$  at the interior pivotal points of the square region shown in the following figure. The values of u at the pivotal points on the boundary are also shown in the figure.

(07 Marks)




Fig. Q7 (c)

8 a. State and prove the recurrence relation of Z – Transformation hence find  $Z_T(n^p)$  and

$$Z_{\rm T} \left[ \cos h \left( \frac{n\pi}{2} + \theta \right) \right]. \tag{07 Marks}$$

b. Find 
$$Z_T^{-1} \left[ \frac{z^3 - 20z}{(z-2)^3 (z-4)} \right]$$
 (06 Marks)

c. Solve the difference equation

$$y_{n+2} - 2y_{n+1} - 3y_n = 3^n + 2n$$
  
Given  $y_0 = y_1 = 0$ . (07 Marks)

\* \* \* \* \*