18MAT11

USN

First Semester B.E. Degree Examination, June/July 2024 Calculus and Linear Algebra

Time: 3 hrs.

Max. Marks: 100

Note: Answer any FIVE full questions, choosing ONE full question from each module.

Module-1

1 a. Find the angle between the curves, $r = a(1 + \sin \theta)$ and $r = a(1 - \sin \theta)$. (08 Marks)

b. Show that for the curve $r(1 - \cos\theta) = 2a$, ρ^2 varies as r^3 . (06 Marks)

c. Derive $\tan \phi = r \frac{d\theta}{dr}$ with usual notation. (06 Marks)

OR

2 a. Find the pedal equation of the curve $r^2 = a^2[\cos 2\theta + \sin 2\theta]$. (08 Marks)

b. If ρ be the radius of curvature at any point P(x, y) on $y^2 = 4ax$, show that $a\rho^2 = 4(x + a)^3$.

(06 Marks)

c. Show that the evolute of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is $(xa)^{2/3} + (yb)^{2/3} = (a^2 - b^2)^{2/3}$. (06 Marks)

Module-2

3 a. Expand $log[1 + e^x]$ using Maclaurin's series upto the term containing x^3 . (06 Marks)

b. Evaluate: $x \to 0 \left[\frac{3^x + 4^x + 5^x}{3} \right]^{1/x}$. (07 Marks)

c. If z = f(x, y) with $x = r \cos \theta$ and $y = r \sin \theta$ show that:

$$\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 = \left(\frac{\partial z}{\partial r}\right)^2 + \frac{1}{r^2} \left(\frac{\partial z}{\partial \theta}\right)^2.$$
 (07 Marks)

OR

4 a. If U = f(x - y, y - z, z - x) prove that $\frac{\partial U}{\partial x} + \frac{\partial U}{\partial y} + \frac{\partial U}{\partial z} = 0$. (06 Marks)

b. Find the Jacobian, $\frac{\partial(x, y, z)}{\partial(r, \theta, \phi)}$ from $x = r\cos\theta\cos\phi$, $y = r\cos\theta\sin\phi$ and $z = r\sin\theta$. (07 Marks)

c. Show that the rectangular box of maximum volume and a given surface area is cube.

(07 Marks)

18MAT11

Module-3

- 5 a. Evaluate: $\iint_{R} x^2 y dx dy$, over the region bounded by the curves $y = x^2$ and y = x. (06 Marks)
 - b. Find the volume generated by the revolution of the cardioide $r = a(1 + \cos \theta)$ about the initial line, using double integral. (07 Marks)
 - c. Using definition of Gamma function, show that $\Gamma(\frac{1}{2}) = \sqrt{\pi}$. (07 Marks)

OR

- 6 a. Evaluate: $\int_{-a}^{a} \int_{0}^{\sqrt{a^2-x^2}} \sqrt{x^2+y^2} \, dy dx$ by changing into polar co-ordinates. (06 Marks)
 - b. Evaluate: $\int_{0}^{1} \int_{0}^{\sqrt{1-x^2}} \int_{0}^{\sqrt{1-x^2-y^2}} xyz \, dz \, dy \, dx.$ (07 Marks)
 - c. Show that $: \int_0^\infty \sqrt{x} e^{-x^2} dx \times \int_0^\infty \frac{e^{-x^2}}{\sqrt{x}} dx = \frac{\pi}{2\sqrt{2}}.$ (07 Marks)

Module-4

- 7 a. A body in air at 25°C cools from 100°C to 75°C in 1 minute. Find the temperature of the body at the end of 3 minutes. (06 Marks)
 - b. Find the orthogonal trajectories of the family of curves $r^n = a^n \sin n\theta$, where 'a' is a parameter. (07 Marks)

c. Solve:
$$y \left(\frac{dy}{dx}\right)^2 + (x-y)\frac{dy}{dx} - x = 0, p = \frac{dy}{dx}$$
. (07 Marks)

OR

- 8 a. Solve: $(2x^2 6xy)dy + (8xy 9y^2)dx = 0$. (06 Marks)
 - b. Solve: $\frac{dy}{dx} + y^2 \tan x = y^3 \sec x.$ (07 Marks)
 - c. The current 'i' in an electrical circuit containing an inductance L and a resistance R in series and, acted upon an e.m.f E sin ot satisfies the differential equation:

$$L\frac{di}{dt} + R.i = E \sin \omega t$$

Find the value of the current at any time, if initially there is no current in the circuit.

(07 Marks)

Module-5

9 a. By applying elementary row operations find rank of:

 $\begin{bmatrix}
1 & 2 & 2 & 1 \\
2 & 1 & 1 & 2 \\
3 & 2 & 2 & 3 \\
0 & 1 & 1 & 0
\end{bmatrix}$ (06 Marks)

b. Solve using Gauss – Seidel method [carry out 4 iterations]:

6x + 15y + 2z = 72 27x + 6y - z = 85x + y + 54z = 110. (07 Marks)

c. Diagonalize the square matrix $A = \begin{bmatrix} -19 & 7 \\ -42 & 16 \end{bmatrix}$. (07 Marks)

OR

10 a. Apply Gauss – Jordan method to solve the following system of equations:

2x + y + 3z = 1 4x + 4y + 7z = 12x + 5y + 9z = 3. (06 Marks)

b. Test for consistency, if consistent solve it.

x + 2y + 3z = 14 4x + 5y + 7z = 353x + 3y + 4z = 21. (07 Marks)

c. Using Rayleigh's power method, find largest eigen value and the corresponding eigen vector of:

 $\begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix}$

By taking $X^{(0)} = [1, 1, 1]^T$ as initial eigen vector. (07 Marks)

* * * * *