

# CBCS SCHEME

USN 

|  |  |  |  |  |  |  |  |
|--|--|--|--|--|--|--|--|
|  |  |  |  |  |  |  |  |
|--|--|--|--|--|--|--|--|

18MAT31

## Third Semester B.E. Degree Examination, Aug./Sept.2020 Transform Calculus, Fourier Series and Numerical Techniques

Time: 3 hrs.

Max. Marks: 100

**Note:** Answer any FIVE full questions, choosing ONE full question from each module.

### Module-1

- 1 a. Find  $L\{e^{-2t}t \cos 2t\}$ . (06 Marks)  
 b. Express the function in terms of unit step function and hence find Laplace transform of :  

$$f(t) = \begin{cases} 1 & 0 \leq t \leq 1 \\ t & 1 < t \leq 2 \\ t^2 & t > 2 \end{cases} \quad (07 \text{ Marks})$$
  
 c. Solve the equation  $y''(t) + 3y'(t) + 2y(t) = 0$  under the condition  $y(0) = 1, y'(0) = 0$ . (07 Marks)

OR

- 2 a. Find :  
 i)  $L^{-1}\left\{\frac{s+3}{s^2-4s+13}\right\}$  ii)  $L^{-1}\left\{\log\frac{(s^2+1)}{s(s+1)}\right\}$  (06 Marks)  
 b. Find  $L^{-1}\left\{\frac{s^2}{(s^2+a^2)^2}\right\}$  using convolution theorem. (07 Marks)  
 c. A periodic function of period  $2a$  is defined by  

$$f(t) = \begin{cases} E & 0 \leq t \leq a \\ -E & a < t \leq 2a \end{cases}$$
  
 Where  $E$  is a constant and show that  $\text{trim } L\{f(t)\} = \frac{E}{S} \tan h\left(\frac{as}{2}\right)$ . (07 Marks)

### Module-2

- 3 a. Express  $f(x) = x^2$  as a Fourier series in the interval  $-\pi < x < \pi$ . Hence deduce that  

$$\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} - \dots = \frac{\pi^2}{12}.$$
 (07 Marks)  
 b. Obtain the Fourier series expression of  $f(x) = \begin{cases} \pi x & 0 < x < 1 \\ \pi(2-x) & 1 < x < 2 \end{cases}$ . (07 Marks)  
 c. Obtain the half range cosine series for the function  $f(x) = (x-1)^2$   $0 \leq x \leq 1$ . (06 Marks)

OR

- 4 a. Obtain the Fourier series of  $f(x) = \begin{cases} \frac{\pi-x}{2}, & 0 < x < 2\pi \\ 0, & \text{elsewhere} \end{cases}$ . Hence deduce that

$$1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots = \frac{\pi}{4}.$$

(07 Marks)

- b. Obtain the half range cosine series of  $f(x) = x \sin x$ ,  $0 \leq x \leq \pi$ .

(07 Marks)

- c. Express  $f(x)$  as a Fourier series upto first harmonic.

|      |   |   |    |   |   |   |
|------|---|---|----|---|---|---|
| x    | 0 | 1 | 2  | 3 | 4 | 5 |
| f(x) | 4 | 8 | 15 | 7 | 6 | 2 |

(06 Marks)

Module-3

- 5 a. Find the Fourier cosine transform of

$$f(x) = \begin{cases} x & \text{for } 0 < x < 1 \\ 2-x & \text{for } 1 < x < 2 \\ 0 & \text{for } x > 2 \end{cases}$$

(07 Marks)

- b. Find the Fourier transform by  $f(x) = e^{-|x|}$ .

(07 Marks)

- c. Obtain the inverse Z - transform by  $u(z) = \frac{z}{(z-2)(z-3)}$ .

(06 Marks)

OR

- 6 a. Find the Fourier transform by

$$f(x) = \begin{cases} 1-|x| & |x| < 1 \\ 0 & |x| > 1 \end{cases}$$

(07 Marks)

and show that  $\int_0^\infty \frac{\sin^2 t}{t^2} dt = \frac{\pi}{2}$ .

- b. Find the z-transform of: i)  $\cos n\theta$  ii)  $\sin n\theta$ .

(06 Marks)

- c. Solve using Z - transform  $u_{n+2} - 4u_n = 0$  given that  $u_0 = 0$  and  $u_1 = 2$ .

(07 Marks)

Module-4

- 7 a. Using Taylor's series method solve  $y(x) = x + y$ ,  $y(0) = 1$  then find  $y$  at  $x = 0.1, 0.2$  consider upto 4<sup>th</sup> degree.

(07 Marks)

- b. Solve  $y'(x) = 1 + \frac{y}{x}$ ,  $y(1) = 2$  then find  $y(1.2)$  with  $n = 0.2$  using modified Euler's method.

(06 Marks)

- c. Solve  $y'(x) = x - y^2$  and the data is  $y(0) = 0$ ,  $y(0.2) = 0.02$ ,  $y(0.4) = 0.0795$ ,  $y(0.6) = 0.1762$  then find  $y(0.8)$  by applying Milne's method and applying corrector formula twice.

(07 Marks)

OR

- 8 a. Solve  $y'(x) = 3x + \frac{y}{2}$ ,  $y(0) = 1$  then find  $y(0.2)$  with  $n = 0.2$  using modified Euler's method. (06 Marks)
- b. Solve  $y(x) = 3e^x + 2y$ ,  $y(0) = 0$  then find  $y(0.1)$  with  $h = 0.1$  using Runge-Kutta method of fourth order. (07 Marks)
- c. Solve  $y'(x) = 2e^x - y$  and data is

|   |   |       |       |       |
|---|---|-------|-------|-------|
| x | 0 | 0.1   | 0.2   | 0.3   |
| y | 2 | 2.010 | 2.040 | 2.090 |

Then find  $y(0.4)$  by using Adam's Bash forth method. (07 Marks)

Module-5

- 9 a. By applying Milne's predictor and corrector method to compute  $y(0.4)$  give the differential equation  $\frac{d^2y}{dx^2} = 1 - \frac{dy}{dx}$  and the following table by initial value. (07 Marks)

|      |   |        |        |        |
|------|---|--------|--------|--------|
| x    | 0 | 0.1    | 0.2    | 0.3    |
| y    | 1 | 1.1103 | 1.2427 | 1.3990 |
| $y'$ | 1 | 1.2103 | 1.4427 | 1.6990 |

- b. Derive Euler's equation in the standard form  $\frac{\partial f}{\partial y} - \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) = 0$ . (06 Marks)
- c. Find the extremal of the functional  $\int_{x_1}^{x_2} (y' + x^2 y'^2) dx$ . (07 Marks)

OR

- 10 a. By Runge Kutta method solve  $\frac{d^2y}{dx^2} = x \left( \frac{dy}{dx} \right)^2 - y^2$  for  $x = 0.2$  correct to four decimal places. (07 Marks)
- Using initial condition  $y(0) = 1$ ,  $y'(0) = 0$ .
- b. Prove that the shortest distance between two points in a plane is a straight line. (06 Marks)
- c. Find the curve on which the functional  $\int_0^1 [y'^2 + 12xy] dx$  with  $y(0) = 0$ ,  $y(1) = 1$ . (07 Marks)

\* \* \* \* \*